



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Bc

Search PubMed for Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Sort Save Text Clip Add Order

Text Version

1: Diabetologia 1992 Jan;35(1):12-8

Related Articles, Links

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Angiotensin converting enzyme inhibition prevents development of muscle and nerve dysfunction and stimulates angiogenesis in streptozotocin-diabetic rats.

Cameron NE, Cotter MA, Robertson S.

School of Biomedical Sciences, University of Aberdeen, Scotland, UK.

The effects of the angiotensin converting enzyme inhibitor lisinopril on slow and fast twitch muscle contractile properties, nerve conduction and hypoxic resistance, and muscle and nerve capillary density were examined in streptozotocin-diabetic rats. Prolongation of soleus contraction and relaxation were partially prevented by treatment (p less than 0.01). A 22% deficit in fast twitch extensor digitorum longus tetanic tension production was also ameliorated (p less than 0.01). Sciatic motor and sensory conduction velocity, 25% and 12% reduced by diabetes respectively, were 75% normalized by lisinopril (p less than 0.01). There was a 47% increase in resistance to hypoxic conduction block with diabetes (p less than 0.01). Lisinopril treatment resulted in normal hypoxic resistance. Capillarization of nerve and muscle was little affected by diabetes; however, there was a 17% increase in capillary density in sciatic nerve, and a 40% increase in extensor digitorum longus muscle with lisinopril (p less than 0.01). For soleus, a smaller treatment-induced increase in capillary density led to an elevated capillary/muscle fibre ratio (p less than 0.01). These results suggest that lisinopril promoted angiogenesis. It was concluded that the beneficial effect of preventive lisinopril treatment is likely to depend upon a reduction of peripheral vascular resistance and improvement of tissue blood flow, which implicates relative hypoxia as an important factor in the development of myopathy and neuropathy in experimental diabetes.

PMID: 1371757 [PubMed - indexed for MEDLINE]

Display Abstract Sort Save Text Clip Add Order

Write to the Help Desk
NCBI | NLM | NIH